



Phenolipids as antioxidants in omega-3 enriched food products

Sørensen, Ann-Dorit Moltke; Aleman, M.; Durand, E.; Villeneuve, P.; Bou, R.; Guardiola, F.; Jacobsen, Charlotte

Publication date:
2013

Document Version
Publisher's PDF, also known as Version of record

[Link back to DTU Orbit](#)

Citation (APA):
Sørensen, A-D. M., Aleman, M., Durand, E., Villeneuve, P., Bou, R., Guardiola, F., & Jacobsen, C. (2013). *Phenolipids as antioxidants in omega-3 enriched food products*. Abstract from 104th AOCS Annual Meeting & Expo, Montreal, Canada.

General rights

Copyright and moral rights for the publications made accessible in the public portal are retained by the authors and/or other copyright owners and it is a condition of accessing publications that users recognise and abide by the legal requirements associated with these rights.

- Users may download and print one copy of any publication from the public portal for the purpose of private study or research.
- You may not further distribute the material or use it for any profit-making activity or commercial gain
- You may freely distribute the URL identifying the publication in the public portal

If you believe that this document breaches copyright please contact us providing details, and we will remove access to the work immediately and investigate your claim.

Phenolipids as antioxidant in omega-3 enriched food products

Ann-Dorit Moltke Sørensen, Mercedes Alemán, Erwann Durand, Pierre Villeneuve, Ricard Bou, Francesc Guardiola & Charlotte Jacobsen

Foods containing omega-3 PUFA are highly susceptible to oxidation. This causes formation of undesirable flavors and loss of health-beneficial fatty acids. To protect these food products, antioxidant addition may be a solution. Lately, extensive work has been performed on phenolipids and their efficacy in model emulsion systems. Since the polar paradox hypothesis was a simplified statement of the antioxidant efficacy in emulsions, a new term, “cut-off effect”, was introduced. The cut-off effect describes the efficacy of phenolipids in simple emulsions. However, most food products consist of a complex matrix where several factors may influence the oxidative stability, e.g. type and concentration of emulsifier. Thus, a better understanding of the antioxidative effect of phenolipids in complex foods is of great interest.

The aim of this study was to evaluate the antioxidative effect of caffeic acid and its esters, caffeates, in two different fish-oil-enriched food products: mayonnaise and milk. Lipid oxidation was evaluated from 3 parameters measured over storage time: peroxide value, volatiles and tocopherol concentrations. The results demonstrate the influence of the complex emulsions on the antioxidant efficacy.